

Labs and Take-Home Prescriptions – 4/11/2017, Planned

Labs and Take-Home Prescriptions – Planned for 4/11/2017

Other

Communication to Provider

Referrals for authorization of drug treatments from the patient's insurance are now being request automatically for both Treatment and Therapy Plans. Most insurances require a minimum of 72-hours to approve drug treatments.

Labs

CBC AND DIFFERENTIAL

Routine, Lab Collect, Expires-S+365, Interval-See Treatment Plan, Count-16

COMPREHENSIVE METABOLIC PANEL

Routine, Lab Collect, Expires-S+365, Interval-See Treatment Plan, Count-16

Research

Research Lab Order

Expires-S+28, Routine, Clinic Collect
Pre-Study Research Labs
Optional: Peripheral Blood Banking (One 10 ml EDTA purple-top)

Research Lab Order

Expires-S+365, Routine, Clinic Collect
First Progression: Optional Peripheral Blood Banking (One 10 ml EDTA purple-top)

Research Lab Order

Expires-S+365, Routine, Clinic Collect
At Progression for Responders: Fresh Biopsy (FFPE block or 12 unstained slides)

Cycle 1 – 4/12/2017 through 5/2/2017 (21 days), Planned

Day 1, Cycle 1 – Planned for 4/12/2017

Other

POCT URINE PREGNANCY

Order details

Nursing Orders

Nurse Verification

I verify the treatment plan.

Nurse Verification

I verify the treatment plan.

Labs needed today

CBC with differential and CMP are needed today.

Research

Research Requirements and Information

Day 1: Physical exam, weight, vital signs ECOG, and concomitant medications are needed.

On days of clinic visits when PK samples are to be drawn, Talazoparib should be taken at the clinic after completion of pre-dose sampling and assessment

Talazoparib capsules whole and not to chew prior to swallowing. Talazoparib can be taken with or without food. No capsule should be ingested if it is broken, cracked or otherwise not intact. Patients should be encouraged to take their dose at approximately the same time each day.

Research (Central) Lab Order, Pre-Dose

Expires-S+1, Routine, Clinic Collect
Pre-dose: PK (One 7 ml K3 EDTA purple-top)

Research (Central) Labs, Post-Dose

▼ Actions ✕

Expires-S+1, Routine, Clinic Collect
Post-dose:

30 mins post dose: PK (One 7 ml K3 EDTA purple-top)


2 hr after first post dose sample: PK (One 7 ml K3 EDTA purple-top)

Chemotherapy

▼ Actions ✕

Investigational Take-Home Medication 1,000 mcg

▼ Actions ✕

 1,000 mcg, Oral, ONCE, Starting at treatment start time, For 1 dose
HIC 1405014058 Talazoparib. Take 1 capsule (1000mg) by mouth once daily for 21 days. Swallow the capsules whole and do not chew prior to swallowing. Take at approximately the same time each day, preferable in the morning, with or without food. Quantity dispensed =30 capsules.

Day 8, Cycle 1 – Planned for 4/19/2017

▼ Actions ✕

Nursing Orders

▼ Actions ✕

Labs needed today

▼ Actions ✕

CBC with differential and CMP are needed today.

Day 15, Cycle 1 – Planned for 4/26/2017

▼ Actions ✕

Nursing Orders

▼ Actions ✕

Labs needed today

▼ Actions ✕

CBC with differential and CMP are needed today.

Cycle 2 – 5/3/2017 through 5/23/2017 (21 days), Planned

▼ Actions ✕

Day 1, Cycle 2 – Planned for 5/3/2017

▼ Actions ✕

Other

▼ Actions ✕

POCT URINE PREGNANCY

▼ Actions ✕

[Order details](#)

Nursing Orders

▼ Actions ✕

Nurse Verification

▼ Actions ✕

[I verify the treatment plan.](#)

Nurse Verification

▼ Actions ✕

[I verify the treatment plan.](#)

Labs needed today

▼ Actions ✕

CBC with differential and CMP are needed today.

Research

▼ Actions ✕

Research Requirements and Information

▼ Actions ✕

Day 1: Physical exam, weight, vital signs ECOG, and concomitant medications are needed.

On days of clinic visits when PK samples are to be drawn, Talazoparib should be taken at the clinic after completion of pre-dose sampling and assessment

Talazoparib capsules whole and not to chew prior to swallowing. Talazoparib can be taken with or without food. No capsule should be ingested if it is broken, cracked or otherwise not intact. Patients should be encouraged to take their dose at approximately the same time each day.

Research (Central) Lab Order, Pre-Dose

▼ Actions ✕

Expires-S+1, Routine, Clinic Collect
Pre-dose:

Optional Peripheral Blood Banking (One 10 ml EDTA purple-top)
PK (One 7 ml K3 EDTA purple-top)

Research (Central) Labs, Post-Dose

Actions X

Expires-S+1, Routine, Clinic Collect
Post-dose:

30 mins post dose: PK (One 7 ml K3 EDTA purple-top)


2 hr after first post dose sample: PK (One 7 ml K3 EDTA purple-top)

Chemotherapy

Actions X

Investigational Take-Home Medication 1,000 mcg

Actions X

 1,000 mcg, Oral, ONCE, Starting at treatment start time, For 1 dose
HIC 1405014058 Talazoparib. Take 1 capsule (1000mg) by mouth once daily for 21 days. Swallow the capsules whole and do not chew prior to swallowing. Take at approximately the same time each day, preferable in the morning, with or without food. Quantity dispensed =30 capsules.

Cycle 3 – 5/24/2017 through 6/13/2017 (21 days), Planned

Actions X

Day 1, Cycle 3 – Planned for 5/24/2017

Actions X

Other

Actions X

POCT URINE PREGNANCY

Actions X

[Order details](#)

Nursing Orders

Actions X

Nurse Verification

Actions X

[I verify the treatment plan.](#)

Nurse Verification

Actions X

[I verify the treatment plan.](#)

Labs needed today

Actions X

CBC with differential and CMP are needed today.

Research

Actions X

Research Requirements and Information

Actions X

Day 1: Physical exam, weight, vital signs ECOG, and concomitant medications are needed.

On days of clinic visits when PK samples are to be drawn, Talazoparib should be taken at the clinic after completion of pre-dose sampling and assessment

Talazoparib capsules whole and not to chew prior to swallowing. Talazoparib can be taken with or without food. No capsule should be ingested if it is broken, cracked or otherwise not intact. Patients should be encouraged to take their dose at approximately the same time each day.

Research (Central) Lab Order, Pre-Dose

Actions X

Expires-S+1, Routine, Clinic Collect
Pre-dose:
Optional Peripheral Blood Banking (One 10 ml EDTA purple-top)
PK (One 7 ml K3 EDTA purple-top)

Research (Central) Labs, Post-Dose

Actions X

Expires-S+1, Routine, Clinic Collect
Post-dose:

30 mins post dose: PK (One 7 ml K3 EDTA purple-top)


2 hr after first post dose sample: PK (One 7 ml K3 EDTA purple-top)

Chemotherapy

Actions X

Investigational Take-Home Medication 1,000 mcg

Actions X

 1,000 mcg, Oral, ONCE, Starting at treatment start time, For 1 dose
HIC 1405014058 Talazoparib. Take 1 capsule (1000mg) by mouth once daily for 21 days. Swallow the capsules whole and do not chew prior to swallowing. Take at approximately the same time each day, preferable in the morning, with or without food. Quantity dispensed =30 capsules.

Cycle 4 – 6/14/2017 through 7/4/2017 (21 days), Planned

Actions X

Day 1, Cycle 4 – Planned for 6/14/2017

Actions X

Other

Actions X

POCT URINE PREGNANCY

Actions X

[Order details](#)

Nursing Orders

Actions X

Nurse Verification

Actions X

[I verify the treatment plan.](#)

Nurse Verification

Actions X

[I verify the treatment plan.](#)

Labs needed today

Actions X

CBC with differential and CMP are needed today.

Research

Actions X

Research Requirements and Information

Actions X

Day 1: Physical exam, weight, vital signs ECOG, and concomitant medications are needed.

On days of clinic visits when PK samples are to be drawn, Talazoparib should be taken at the clinic after completion of pre-dose sampling and assessment

Talazoparib capsules whole and not to chew prior to swallowing. Talazoparib can be taken with or without food. No capsule should be ingested if it is broken, cracked or otherwise not intact. Patients should be encouraged to take their dose at approximately the same time each day.

Research (Central) Lab Order, Pre-Dose

Actions X

Expires-S+1, Routine, Clinic Collect

Pre-dose:

Optional Peripheral Blood Banking (One 10 ml EDTA purple-top)

PK (One 7 ml K3 EDTA purple-top)

Research (Central) Labs, Post-Dose

Actions X

Expires-S+1, Routine, Clinic Collect

Post-dose:

30 mins post dose: PK (One 7 ml K3 EDTA purple-top)

2 hr after first post dose sample: PK (One 7 ml K3 EDTA purple-top)

Chemotherapy

Actions X

Investigational Take-Home Medication 1,000 mcg

Actions X

 1,000 mcg, Oral, ONCE, Starting at treatment start time, For 1 dose

HIC 1405014058 Talazoparib. Take 1 capsule (1000mg) by mouth once daily for 21 days. Swallow the capsules whole and do not chew prior to swallowing. Take at approximately the same time each day, preferable in the morning, with or without food. Quantity dispensed =30 capsules.

Cycle 5 – 7/5/2017 through 7/25/2017 (21 days), Planned

Actions X

Day 1, Cycle 5 – Planned for 7/5/2017

Actions X

Other

Actions X

POCT URINE PREGNANCY

Actions X

[Order details](#)

Nursing Orders

Actions X

Nurse Verification

Actions X


[I verify the treatment plan.](#)

Nurse Verification

Actions X

I verify the treatment plan.

Labs needed today

▼ Actions 

CBC with differential and CMP are needed today.

Research

▼ Actions 

Research Requirements and Information

▼ Actions 

Day 1: Physical exam, weight, vital signs ECOG, and concomitant medications are needed.


On days of clinic visits when PK samples are to be drawn, Talazoparib should be taken at the clinic after completion of pre-dose sampling and assessment


Talazoparib capsules whole and not to chew prior to swallowing. Talazoparib can be taken with or without food. No capsule should be ingested if it is broken, cracked or otherwise not intact. Patients should be encouraged to take their dose at approximately the same time each day.

Chemotherapy

▼ Actions 

Investigational Take-Home Medication 1,000 mcg

▼ Actions 

 1,000 mcg, Oral, ONCE, Starting at treatment start time, For 1 dose
HIC 1405014058 Talazoparib. Take 1 capsule (1000mg) by mouth once daily for 21 days. Swallow the capsules whole and do not chew prior to swallowing. Take at approximately the same time each day, preferable in the morning, with or without food. Quantity dispensed =30 capsules.

Cycle 6 – 7/26/2017 through 8/15/2017 (21 days), Planned

▼ Actions 

Cycle 7 – 8/16/2017 through 9/5/2017 (21 days), Planned

▼ Actions 

Cycle 8 – 9/6/2017 through 9/26/2017 (21 days), Planned

▼ Actions 

Cycle 9 – 9/27/2017 through 10/17/2017 (21 days), Planned

▼ Actions 

Cycle 10 – 10/18/2017 through 11/7/2017 (21 days), Planned

▼ Actions 

Cycle 11 – 11/8/2017 through 11/28/2017 (21 days), Planned

▼ Actions 

Cycle 12 – 11/29/2017 through 12/19/2017 (21 days), Planned

▼ Actions 

Cycle 13 – 12/20/2017 through 1/9/2018 (21 days), Planned

▼ Actions 

Cycle 14 – 1/10/2018 through 1/30/2018 (21 days), Planned

▼ Actions 

Cycle 15 – 1/31/2018 through 2/20/2018 (21 days), Planned

▼ Actions 