

Table 1. Baseline Characteristics

| | PCPT (n=18,880) | |
|--|-----------------|-------|
| | N | % |
| Age at enrollment (median [IQR],years) | 63 [59-67] | |
| 50-54 | 3 | 0.0% |
| 55-59 | 5,908 | 31.3% |
| 60-64 | 5,794 | 30.7% |
| ≥65 | 7,175 | 38.0% |
| Follow-up for those last known to be alive (median [IQR],years) | 79 [76-83] | |
| < 5 | 5 | 0.0% |
| 5.1 – 10 | 292 | 2.3% |
| 10.1 – 15.0 | 106 | 0.8% |
| >15 | 12,432 | 96.7% |
| Race/Ethnicity | | |
| Non-Hispanic white | 17,382 | 92.1% |
| Non-Hispanic African American | 712 | 3.8% |
| Hispanic | 493 | 2.6% |
| Other | 293 | 1.6% |
| History of prostate cancer in 1 st degree relative | | |
| Yes | 2,913 | 15.4% |
| PSA at entry (median [IQR], ng/ml) | 1.1 [0.7-1.7] | |
| 0 – 1.0 | 9,120 | 48.3% |
| 1.1 – 2.0 | 6,715 | 35.6% |
| 2.1 – 3.0 | 3,044 | 16.1% |
| 3.1 – 4.0 | 1 | 0.0% |
| Education (highest level) | | |
| ≤ High school graduate or GED | 3,569 | 18.9% |
| Some college/vocational school | 5,509 | 29.2% |
| ≥ College graduate | 9,790 | 51.9% |
| Cigarette smoking | | |
| Never | 6,218 | 32.9% |
| Former | 11,171 | 59.2% |
| Current | 1,486 | 7.9% |
| Body mass index (kg/m ²) | | |
| <25 | 4,817 | 25.5% |
| 25-<30 | 9,499 | 50.3% |
| ≥30 | 4,378 | 23.2% |
| Numbers for each descriptive factor which are unknown are not presented. | | |